

Negative Pressure Wound Therapy and the Perfusion Paradox: Reconciling Clinical Success with Foundational Principles of Tissue Healing

Opinion

Volume 2 Issue 1- 2026

Author Details

Colton C Mowers¹, Michael S Shuler², Brett A Freedman^{3*}

¹Department of Orthopaedic Surgery, Rush Medical College, USA

²Department of Orthopedic Surgery, Athens Orthopedic Clinic, USA

³Department of Orthopaedic Surgery, Mayo Clinic, USA

*Corresponding author

Brett A Freedman, Department of Orthopaedic Surgery, Mayo Clinic, Rochester, MN

Article History

Received: January 28, 2026 Accepted: January 29, 2026 Published: January 30, 2026

Abstract

Negative pressure wound therapy (NPWT) has become a cornerstone of modern wound management because it often reduces wound burden while accelerating closure. Despite this success, a persistent physiologic paradox remains. Negative pressure applied at the wound interface generates compressive forces against the wound bed, impairing tissue perfusion at commonly used therapeutic pressure settings. Traditional foam-based wound fillers require high negative pressure settings (-125mmHg) to achieve effective exudate clearance, narrowing the therapeutic window between exudate management and perfusion preservation. High quality clinical trials across multiple wound types have failed to demonstrate consistent superiority of NPWT over conventional dressings, particularly in high-risk wounds (i.e. dysvascular, pressure-related, infected), where perfusion may be a more significant factor. These conflicting clinical and physiological findings suggest that the limitations of current systems may relate less to the generic concept of NPWT and more to long standing dependence on high-resistance ROCF interfaces. This paradox begs the question, as to whether an optimal wound interface that possesses high outflow efficiency enabling lower therapeutic negative pressure settings could resolve this paradox and improve clinical outcomes, especially in high-risk wounds.

Discussion

Negative pressure wound therapy (NPWT) is commonly described as the application of sub-atmospheric pressure to tissue. In practice, NPWT improves exudate control, reduces dressing change burden, and accelerates wound healing, in many cases, which has driven widespread adoption [1] However, according to Newton's 3rd law of motion, application of negative pressure to a sealed NPWT wound dressing must transmit an opposing force against the wound bed, resulting in compression of the wound surface [2,3].

Low magnitude compression (20-30mmHg) may be beneficial by reducing interstitial edema, improving lymphatic drainage and improving local perfusion. However, compression has dose-dependent effect on tissue healing, and increasing compressive forces, especially when continuously applied for days at a time, will eventually impair microvascular perfusion, particularly in dysvascular tissue or wounds over hard prominences [2]. Perfusion studies demonstrate progressive reductions in blood flow beneath negative pressure dressings as suction

magnitude increases, challenging the assumption that negative pressure inherently improves perfusion [2-4].

Despite these physiological findings, NPWT systems have relied almost exclusively on reticulated open cell foam (ROCF) as the wound filler of choice. This material, characterized by pore sizes of approximately 400 to 600 microns, exhibits increasing resistance to flow under compression [2,3]. As resistance increases, higher negative pressure settings are required to achieve effective fluid evacuation. Experimental work demonstrates that maximal exudate removal with foam-based interfaces occur around -125mmHg, even though maximal wound contraction occurs at lower pressures [4]. This material-specific requirement for high negative pressure settings creates a narrow therapeutic window for ROCF in which clinicians must balance fluid management with perfusion preservation.

The mechanisms underlying the therapeutic effect of NPWT remain incompletely defined despite decades of use. Proposed therapeutic effects include macrodeformation, microdeformation, and altered cellu-

lar signaling [5]. The paradox between commonly observed favorable clinical outcomes with less dressing maintenance requirements when using NPWT with ROCF compared to the adverse effect this particular approach has on wound bed perfusion has been known for quite some time, but under-appreciated, possibly due to the lack of an effective alternative NPWT interface. This has fostered confirmation bias, whereby favorable clinician experiences overshadow contradictory physiologic data and inconsistent clinical trials [6].

High quality clinical trials demonstrate a lack of consistent patient centered benefits despite widespread clinical adoption of NPWT with ROCF. Randomized studies in open fractures, incisional wounds after major extremity trauma, and surgical wounds healing by secondary intention have failed to demonstrate meaningful improvements in patient centered outcomes compared with standard dressings [7-9]. Systematic reviews similarly fail to consistently find superior heal-

ing with NPWT, especially in high-risk wound categories [6,10]. The SAWHI trial in subcutaneous abdominal wound healing impairment showed no improvement in healing with NPWT and reported higher wound-related adverse events compared with conventional care [11]. Across multiple wound categories, higher level evidence shows inconsistent benefit of NPWT and some even signal potential harm in select populations [10,12,13]. Optimizing the wound filler to permit effective therapy at lower pressures may better align clinical performance with basic principles of tissue perfusion and wound healing. It is rare that the first generation of any medical device or key component of a medical device is optimal (Figure 1). However, despite substantial efforts to improve the pumps and adhesive drapes over the last 30+ years, as well as the addition of instillation, the basic material of NPWT remains ROCF. Maybe it's time to consider a change?




Figure 1: AI-generated illustration demonstrating the paradox of NPWT with ROCF. High negative pressure (-125mmHg) is needed to evacuate exudate (black arrow) through ROCF, which is felt to be beneficial to wound healing, but this results in reciprocal positive pressure against the wound (purple arrows) sufficient to cause hypoperfusion of the wound bed.

References

1. Morykwas MJ, Faler BJ, Pearce DJ, Argenta LC (2001) Effects of varying levels of subatmospheric pressure on the rate of granulation tissue formation in experimental wounds in swine. *Ann Plast Surg* 47(5): 547-551.
2. Kairinos N, Voogd AM, Botha PH (2009) Negative-pressure wound therapy II: negative-pressure wound therapy and increased perfusion. Just an illusion? *Plast Reconstr Surg* 123(2): 601-612.
3. Kairinos N, Solomons M, Hudson DA (2009) Negative-pressure wound therapy I: the paradox of negative-pressure wound therapy. *Plast Reconstr Surg* 123(2): 589-598.
4. Borgquist O, Ingemansson R, Malmjö M (2011) The influence of low- and high-pressure levels during negative-pressure wound therapy on wound contraction and fluid evacuation. *Plast Reconstr Surg* 127(2): 551-559.
5. Malmjö M, Gustafsson L, Lindstedt S, Gesslein B, Ingemansson R (2012) The effects of variable, intermittent, and continuous negative pressure wound therapy, using foam or gauze, on wound contraction, granulation tissue formation, and ingrowth into the wound filler. *Eplasty* 12: e5.
6. Ubbink DT, Westerbos SJ, Nelson EA, Vermeulen H (2008) A systematic review of topical negative pressure therapy for acute and chronic wounds. *Br J Surg* 95(6): 685-692.
7. Arundel C, Mandefield L, Fairhurst C (2025) Negative pressure wound therapy versus usual care in patients with surgical wound healing by secondary intention in the UK (SWHSI-2): an open-label, multicentre, parallel-group, randomised controlled trial. *Lancet* 405(10490): 1689-1699.
8. Costa ML, Achten J, Bruce J (2018) Effect of Negative Pressure Wound Therapy vs Standard Wound Management on 12-Month Disability Among Adults With Severe Open Fracture of the Lower Limb. *JAMA* 319(22): 2280-2288.
9. Costa ML, Achten J, Knight R (2020) Effect of Incisional Negative Pressure Wound Therapy vs Standard Wound Dressing on Deep Surgical Site Infection After Surgery for Lower Limb Fractures Associated With Major Trauma: The WHIST Randomized Clinical Trial. *JAMA* 323(6): 519-526.
10. Shi J, Gao Y, Tian J (2023) Negative pressure wound therapy for treating pressure ulcers. *Cochrane Database Syst Rev* 5(5): CD011334.
11. Seidel D, Diedrich S, Herrle F (2020) Negative Pressure Wound Therapy vs Conventional Wound Treatment in Subcutaneous Abdominal Wound Healing Impairment: The SAWHI Randomized Clinical Trial. *JAMA Surg* 155(6): 469-478.
12. Goldman AH, Tetsworth K (2023) AAOS Clinical Practice Guideline Summary: Prevention of Surgical Site Infection After Major Extremity Trauma. *J Am Acad Orthop Surg* 31(1): e1-e8.

